

4

DTIC File Copy

OFFICE OF NAVAL RESEARCH

Contract N00014-86-K-0043

TECHNICAL REPORT No. 111

Optical Nutation in Polymers Irradiated by Ultrashort Laser Pulses

by

Xiao-shen Li, D. L. Lin, Thomas F. George and Xin Sun

Prepared for Publication

in

Physical Review B

Departments of Chemistry and Physics  
State University of New York at Buffalo  
Buffalo, New York 14260

September 1989

Reproduction in whole or in part is permitted for any purpose of the  
United States Government.

This document has been approved for public release and sale;  
its distribution is unlimited.

DTIC  
ELECTE  
SEP 26 1989  
S E D

89 9 25 042

AD-A212 927

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S)  UBUFFALO/DC/89/TR-111			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York		6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260			7b. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER  Contract N00014-86-K-0043		
8c. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.
					WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification)  Optical Nutation in Polymers Irradiated by Ultrashort Laser Pulses					
12. PERSONAL AUTHOR(S) Xiao-shen Li, D. L. Lin, Thomas F. George and Xin Sun					
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) September 1989	
				15. PAGE COUNT 20	
16. SUPPLEMENTARY NOTATION  Prepared for publication in Physical Review B					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	POLYMERS, OPTICAL NUTATION		
			IRRADIATION, PHONON EFFECTS		
			SHORT LASER PULSES, EXCITONS		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) The transient behavior of the optical susceptibility of polydiacetylene induced by an ultrafast pump field is investigated. Within a two-level model which includes phonon effects phenomenologically, an analytical expression for the nonlinear susceptibility is obtained. In addition to spectral hole burning, the novel phenomenon of optical nutation is found. Both this nutation and the shape of the hole depend sensitively on the detuning between the exciton frequency and the sum of the pump field and the phonon mode frequencies. The electronic state and phonon-mediated optical Stark blue shift are also found in this model. The results are in qualitative agreement with experiments and indicate that the steady-state approximation is reliable only when the pulse of the pump field is longer than several exciton lifetimes.					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE (Include Area Code) (202) 696-4410		22c. OFFICE SYMBOL

Optical nutation in polymers irradiated by ultrashort laser pulses

Xiao-shen Li\*, D. L. Lin and Thomas F. George<sup>†</sup>  
Department of Physics and Astronomy  
State University of New York at Buffalo  
Buffalo, New York 14260

and

Xin Sun  
Center of Theoretical Physics  
Chinese Center of Advanced Science and Technology (World Laboratory)  
P. O. Box 8730  
Beijing 100080, People's Republic of China

and

Department of Physics  
Fudan University  
Shanghai 200433, People's Republic of China

Abstract

The transient behavior of the optical susceptibility of polydiacetylene induced by an ultrafast pump field is investigated. Within a two-level model which includes phonon effects phenomenologically, an analytical expression for the nonlinear susceptibility is obtained. In addition to spectral hole burning, the novel phenomenon of optical nutation is found. Both this nutation and the shape of the hole depend sensitively on the detuning between the exciton frequency and the sum of the pump field and the phonon mode frequencies. The electronic state and phonon-mediated optical Stark blue shift are also found in this model. The results are in qualitative agreement with experiments and indicate that the steady-state approximation is reliable only when the pulse of the pump field is longer than several exciton lifetimes.

1988 PACS Nos.: 42.65.-k, 71.35.+z, 36.20.-r

\* On leave of absence from the Shanghai Institute of Metallurgy, Chinese Academy of Sciences, Shanghai, People's Republic of China.

<sup>†</sup> Also affiliated with the Center for Electronic and Electro-optical Materials, State University of New York at Buffalo.



on For

RA&I

B

nced

cation

By

Distribution/

Availability Codes

Dist

Avail and/or

Special

A-1

## I. Introduction

Coherent transient interactions between materials and optical fields are one of the most interesting subjects in physics. Investigations of transient optical effects have become very important in the development of coherent optical spectroscopy and fast response nonlinear optical materials. Such studies cover such a wide spectrum of subjects and materials as photon echo, free induction decay and optical nutation in gaseous materials,<sup>1-5</sup> ultrafast dynamical blue shift of the exciton resonance accompanied by strong bleaching in quantum wells in heterostructure semiconductors,<sup>6-9</sup> nonlinear optical response of excitons<sup>10,11</sup> and large susceptibilities<sup>12,13</sup> in polymers, optical nutation in direct-gap bulk semiconductors,<sup>14,15</sup> and so forth.

A nonlinear optical material can be defined as one whose optical properties change when light shines upon it. In recent years, the study of coherent nonlinear optical processes produced by the excitation of semiconductors in the transparent region well below the absorption edge<sup>6-9</sup> has been extended to polymers.<sup>10,11</sup> When the frequency of the laser beam falling on semiconductors is tuned below the exciton resonance, it is known that virtual excitons are generated. These excitons interact with photons in exactly the same way as real excitons.<sup>7,9-11</sup> Such excitonic effects can be explored more naturally by probing the susceptibility<sup>9</sup> and looking for transient changes that persist as long as the excitation. The imaginary part of the susceptibility is usually examined via the absorption spectrum, and its real part is explored by measuring the index of refraction, provided that the sample is strongly excited well below the conduction band edge.

It is well known that polymers exhibit large nonlinear optical susceptibilities. In comparison with other materials, polymers have the added advantage of having extremely fast ground-state recovery times, which is why

they attract much attention for their potential applications in future generations of signal processing devices. Polydiacetylene (PDA) may be a good candidate for such considerations because of its large  $\chi^{(3)}$  and small transmission loss  $\alpha$  which can be reduced to as low as<sup>16</sup>  $\alpha < 1$  db/cm. Thus the material possesses a fairly large ratio  $\chi^{(3)}/\alpha$ , which is usually the measure of the usefulness of a material employed in switching devices. Besides, PDA is easier to use in the construction of waveguides.<sup>17</sup> In spite of the considerable amount of research carried out so far, however, the mechanism that governs its nonlinear optical response is still not well understood, although tremendous progress has made it ripe for further examination.

In this paper, we study the transient behavior of the induced optical susceptibility of polymers due to irradiation by an ultrashort pulse of a pump laser field. We take, as an example, polydiacetylene-toluene sulfonate (PTS) which is irradiated by a strong pump laser and a weak probing laser. The two lasers may have pulse duration of femtoseconds to picoseconds. For the PTS system we adopt the two-level model<sup>4,6,15</sup> to describe its ground-state and virtual exciton state. The lifetime of the exciton and the mediating effects of the phonon are introduced phenomenologically. We use the perturbation technique to find the differential nonlinear susceptibility by solving analytically to all orders in the pump field and to first order in the test field.

## II. Two-level model

The two electronic states of PTS are denoted by  $|+\rangle$  and  $|-\rangle$ . We assume a dipole transition and define the dipole operators

$$S^- = |-\rangle\langle +| \quad (1a)$$

$$S^+ = |+\times-| \quad , \quad (1b)$$

with the inversion operator given by

$$S^Z = \frac{1}{2} (|+\times+| - |-\times-|) \quad . \quad (1c)$$

For a single virtual exciton and a single phonon mode, we can write the free Hamiltonian as

$$H_0 = \hbar\omega_x S^Z + \hbar\omega_i b_i^\dagger b_i \quad , \quad (2a)$$

where  $b_i^\dagger$  ( $b_i$ ) creates (annihilates) a phonon in mode  $i$  with frequency  $\omega_i$ , and  $\omega_x$  is the exciton frequency.<sup>11</sup> For the interaction Hamiltonian we consider that each virtual exciton interacts with both a strong pump field of amplitude  $E_p$  and frequency  $\omega_p$  and a weak test field of amplitude  $E_t$  and frequency  $\omega_t$ . Since the test field is very weak, we assume the mediating effects of the phonons on the dipole transition to be induced only by the pump field, as shown in Fig. 1. Therefore, we can write the interaction Hamiltonian in both the rotating-wave approximation and the interaction representation as

$$H_{Pi} = - \frac{1}{2} \lambda E_p e^{i\Delta_p t} b_i^\dagger S^- + \text{h.c.} \quad (2b)$$

for the interaction between the pump field and the PTS, and

$$H_t = - \frac{1}{2} \mu E_t e^{i\Delta_t t} S^- + \text{h.c.} \quad (2c)$$

for the interaction between the test field and the PTS, where  $\Delta_{p_i} = \omega_p + \omega_i - \omega_x$ ,  $\Delta_t = \omega_t - \omega_x$ ,  $\mu$  is the dipole matrix element, and  $\lambda$  is a coupling constant proportional to the transition dipole moment  $\mu$ . Here we have neglected the momentum dependence of virtual excitons<sup>11,15</sup> and the damping effect of the phonons. This Hamiltonian uses dipole or projection operators to describe the dipole transition between the ground state and the excitonic state of the material by external fields. It is different from that of Ref. 11, where creation and annihilation operators were used for virtual excitons.

For simplicity, we assume that each phonon mode has only two states,  $|1\rangle$  and  $|0\rangle$ , and that the two and higher-phonon effects are negligible. The initial condition we consider here is that the system electronically is in its ground state with no virtual exciton present, while the phonon mode is in its excited state  $|1\rangle$ . We define  $C_{i\pm}(t)$  as the probability amplitude of the state  $|\pm\rangle$  when there is the  $i$ -th phonon mode only, of which the time derivative  $\dot{C}_{i\pm}$  is composed of three parts: pump field-induced part  $\dot{C}_{i\pm}^p$ , test field-induced part  $\dot{C}_{i\pm}^t$ , and damping part  $\dot{C}_{i\pm}^d$ . By using the Schrödinger equation and the interaction Hamiltonians (2), we obtain<sup>14,15</sup>

$$\dot{C}_{i+}^{p,t}(t) = i\Omega_{p,t}^* e^{-i\Delta_{p_i,t}t} C_{i-}, \quad (3a)$$

$$\dot{C}_{i-}^{p,t}(t) = i\Omega_{p,t} e^{i\Delta_{p_i,t}t} C_{i+}, \quad (3b)$$

where in deriving the equation for  $\dot{C}_{\pm}^t$  we have neglected the phonon effects and have defined  $\Omega_p = \lambda E_p / 2\hbar$  and  $\Omega_t = \mu E_t / 2\hbar$ . Then we may write the damping part phenomenologically<sup>14,15</sup> as

$$\dot{C}_{i+}^d(t) = -\gamma C_{i+} \quad (4a)$$

$$\dot{C}_{i-}^d(t) = 0 \quad (4b)$$

where  $\gamma$  is the damping rate describing the short lifetime of the virtual exciton. Therefore, from  $\dot{C}_{i\pm} = \dot{C}_{i\pm}^p + \dot{C}_{i\pm}^t + \dot{C}_{i\pm}^d$  we have the following equations for  $C_{i\pm}$ :

$$\dot{C}_{i+}(t) = i\Omega_p^* e^{-i\Delta_{pi}t} C_- + i\Omega_t^* e^{-i\Delta_{ti}t} C_{i-} - \gamma C_{i+} \quad (5a)$$

$$\dot{C}_{i-}(t) = i\Omega_p e^{i\Delta_{pi}t} C_{i+} + i\Omega_t e^{i\Delta_{ti}t} C_{i+} \quad (5b)$$

Since the test field is in most cases much weaker than the pump field, we can safely apply perturbation theory to solve (5). The zeroth-order equations are obtained by setting  $E_t = 0$  in (5), namely,

$$\dot{C}_{i+}^o = i\Omega_p^* e^{-i\Delta_{pi}t} C_{i-}^o - \gamma C_{i+}^o \quad (6a)$$

$$\dot{C}_{i-}^o = i\Omega_p e^{i\Delta_{pi}t} C_{i+}^o \quad (6b)$$

In what follows, we assume for simplicity  $\mu = \mu^*$ ,  $\lambda = \lambda^*$  and  $\Omega_{p,t} = \Omega_{p,t}^*$ . With the initial conditions  $C_{i+}^o(0) = 0$  and  $C_{i-}^o(0) = 1$ , we find the solutions

$$C_{i+}^o = \frac{i\Omega_p}{\Omega_i} \sin\Omega_i t \exp[-(i\Delta_{pi} + \gamma)t/2] \quad (7a)$$



$$C_{i-}^0 = (\cos \Omega t + \frac{\gamma - i\Delta_{p_i}}{2\Omega_i} \sin \Omega_i t) e^{(i\Delta_{p_i} - \gamma)t/2}, \quad (7b)$$

where we have defined the complex Rabi frequency

$$\Omega_i = \frac{1}{2} [(\Delta_{p_i} + i\gamma)^2 + 4\Omega_p^2]^{1/2}. \quad (8)$$

The real part of the Rabi frequency is given by

$$\Omega_R = [(\sqrt{x^2 + y^2} + x)/8]^{1/2}, \quad (9a)$$

which determines the oscillation frequency, and the imaginary part by

$$\Omega_I = -[(\sqrt{x^2 + y^2} - x)/8]^{1/2}, \quad (9b)$$

which determines the envelope of the oscillation. In (9) we have defined

$$x = \Delta_{p_i}^2 + 4\Omega_p^2 - \gamma^2 \text{ and } y = 2\Delta_{p_i}\gamma, \text{ with } \Delta_{p_i} < 0.$$

The solutions up to first-order perturbation can be obtained by inserting the zeroth-order solutions (7) into the right-hand side of (5). The results are

$$C_{i\pm}(t) = C_{i\pm}^0(t) + \delta C_{i\pm}(t) + A_{i\pm} \quad (10)$$

$$\delta C_{i-} = \frac{\Omega_t \Omega_p}{\Omega_i} \frac{\exp[i\Delta_t t - \frac{1}{2}(i\Delta_{p_i} + \gamma)t]}{[\frac{1}{2}(i\Delta_{p_i} + \gamma) - i\Delta_t]^2 + \Omega_i^2}$$

$$\times \left[ \left( \frac{1}{2}(i\Delta_{p_i} + \gamma) - i\Delta_t \right) \sin\Omega_i t + \Omega_i \cos\Omega_i t \right] \quad (11a)$$

$$\begin{aligned} \delta C_{i+} = & \frac{i\Omega_t}{2\Omega_i} \frac{\exp[-i\Delta_t t + \frac{1}{2}(i\Delta_{p_i} - \gamma)t]}{[i\Delta_t - \frac{1}{2}(i\Delta_{p_i} - \gamma)]^2 + \Omega_i^2} \\ & \times \{[(\Delta_{p_i} + i\gamma)^2 + 2\Omega_p^2 + i\Delta_t(i\Delta_{p_i} - \gamma)] \sin\Omega_i t \\ & - 2\Omega_i[i(\Delta_t - \Delta_{p_i}) + \gamma] \cos\Omega_i t\} . \end{aligned} \quad (11b)$$

The integration constants  $A_{i\pm}$  in (10) are determined by the initial conditions.

The expectation value of the dipole moment of a virtual exciton is

$$\begin{aligned} \mu_i(t) = \mu \langle S_i^-(t) \rangle e^{-i\omega_x t} &= \mu C_{i-}^* C_{i+} e^{-i\omega_x t} \\ &= \mu (C_{i-}^{0*} + \delta C_{i-}^* + A_{i-}^*) (C_{i+}^0 + \delta C_{i+} + A_{i+}) e^{-i\omega_x t} . \end{aligned} \quad (12)$$

It is evident that the dipole moment induced by the test beam is given by

$$\delta \mu_i(t) = \mu (C_{i-}^{0*} \delta C_{i+} + C_{i+}^0 \delta C_{i-}^*) e^{-i\omega_x t} . \quad (13)$$

Therefore, the nonlinear optical susceptibility experienced by the test beam is simply

$$\begin{aligned}
x_i &= n_i \delta \mu_i(t) / (E_t e^{-i\omega_t t}) \\
&= n_i \frac{|\mu|^2}{2\hbar} e^{-\gamma t} \left( \frac{1}{2\Omega_i z_1^i} [\cos \Omega_i^* t + i \frac{\Delta_{p_i} - i\gamma}{2\Omega_i^*} \sin \Omega_i^* t] \right. \\
&\quad \times [(\Delta_{p_i} + i\gamma)^2 + 2\Omega_p^2 - \Delta_t(\Delta_{p_i} + i\gamma)] \sin \Omega_i t \\
&\quad - 2\Omega_i (i(\Delta_t - \Delta_{p_i}) + \gamma) \cos \Omega_i t] \\
&\quad \left. - \frac{\Omega_p^2}{|\Omega_i|^2 z_2^i} \sin \Omega_i t \left[ \left( \frac{1}{2}(i\Delta_{p_i} - \gamma) - i\Delta_t \right) \sin \Omega_i^* t + \Omega_i^* \cos \Omega_i^* t \right] \right) , \quad (14)
\end{aligned}$$

where

$$z_1^i = \left[ \frac{1}{2}(i\Delta_{p_i} - \gamma) - i\Delta_t \right]^2 + \Omega_i^2 \quad (15a)$$

$$z_2^i = \left[ \frac{1}{2}(i\Delta_{p_i} - \gamma) - i\Delta_t \right]^2 + \Omega_i^{*2} . \quad (15b)$$

We have introduced in (14) the optically-induced transient virtual exciton density  $n_i$  which is taken to be proportional to the probability of virtual exciton state,<sup>18</sup>

$$n_i = n_s C_{i+}^o C_{i+}^{o*} = n_s \frac{\Omega_p^2}{|\Omega_i|^2} |\sin \Omega_i t|^2 \exp(-\gamma t) , \quad (16)$$

where we have assumed that all the virtual excitons are induced by the pump field  $E_p$ , since those induced by the weaker probe field are much less likely, and  $n_s$  is the saturation density of the virtual excitons. In this paper we only consider the two phonon modes coupling most strongly to the exciton,<sup>19</sup> so that we have the nonlinear susceptibility

$$\chi_t = \sum_{i=1}^2 \chi_i \quad (17)$$

The transient behavior of the nonlinear optical susceptibility  $\chi_t$  is calculated numerically as a function of time and the frequencies  $\omega_p$  and  $\omega_t$ . The results are presented and discussed in the next section.

### III. Results and discussion

In our numerical study, we have used the unit  $\hbar = 1$ . Other parameters are chosen as<sup>10,11,20</sup>  $\omega_1 = 0.184$  eV,  $\omega_2 = 0.258$  eV,  $\gamma = 0.02$  eV,  $\frac{1}{e}|\mu|^2 n_s = \frac{1}{40\pi}$  eV,  $\Omega_p = 5 \times 10^{-4}$  eV and  $\omega_x = 1.978$  eV. The real and imaginary parts of  $\chi_t$  are computed separately. Figure 2 shows snapshots at different instants of time of the variation of  $\chi_t$  with the pump detuning. The variation with time for fixed detuning is shown in Fig. 3. Figure 4 shows snapshots at different instants of time of the imaginary part of  $\chi_t$  when the resonance condition is satisfied, i.e.,  $\omega_x = \omega_p + \omega_2$ .

It is clearly seen from Fig. 2 that both the differential absorption and refraction are direct effects of the virtual excitons induced by the pump field. The mediating role played by the phonon is substantial when the duration of the pump pulse is long enough (see the two peaks of the imaginary part, where  $\omega_x = \omega_p + \omega_1$ ). This is in agreement with the experimental results

of Ref. 11. However, when the duration of the pump pulse is short, e.g., less than the lifetime of the virtual exciton, the peak structure of the imaginary part is not clear, i.e., we cannot see evident optical phonon-mediated effects. It should be noted at this point that the present results are different from those reported in Ref. 21 in which steady-state susceptibilities are calculated in the three-level two-field model without phonon effects.

Optical nutation can be observed in polymers in the ultrashort time regime when virtual excitons are generated by a pump field tuned below the exciton resonance. This situation can be seen from Fig. 3. However, when the sum of the pump field frequency and one of the phonon frequencies is on resonance with the exciton frequency, the Rabi oscillation amplitude becomes smaller. This reflects the fact that the optical phonon plays a crucial role in the optical nutation of PTS. It is also clearly shown that sometimes the oscillation centers of  $\text{Im}\chi_t$  change with time. This is quite different from nutation phenomena predicted in Refs. 14 and 15 for semiconductors where the oscillation amplitude decreases monotonically like damped oscillators.

From Fig. 4, we can see the transient behavior of the spectral hole burning, which also needs sufficient long duration of the pump pulse to have an effect. The hole is close to the exciton resonance, and there is a deeper hole when  $\omega_x = \omega_p + \omega_i$ . This indicates that there is energy transfer from the pump field to the test field via optical phonons and virtual excitons, also in qualitative agreement with experiments.<sup>10,11</sup>

Since it takes time for the pump field to build up enough virtual excitons and since the virtual excitons are short-lived, their effects are appreciable and stable most of the time during the pulse (on the order of picoseconds). The response time can be anywhere between 20 - 200 femtoseconds

according to our calculation. Both the absorption and refraction parts show up almost right away and taper off rather slowly after reaching their peaks. So, for longer excitation time of the pump field, say, several times of the exciton lifetime, the steady-state method may be reliable.

In addition, we have also found that increasing the intensity of the pump field leads to a deeper and wider hole in the absorption spectrum, which is qualitatively in agreement with the steady-state results of Ref. 11. What seems to be more interesting is that we find a blue shift of 0.005 eV in our calculation. This represents joint effects of both the phonon-induced optical Stark shift and the electronic state optical Stark shift. Such effects have been observed in a recent experiment<sup>20</sup> in which the differential transmission of polydiacetylene-toluene sulfonate has been measured by means of coherent inverse Raman spectroscopy. It is demonstrated in Ref. 20 that this ac Stark effect is important in the determination of the nonlinear optical response of PTS even in the small signal limit. The effect is reproduced theoretically in Ref. 20 by including an extra term to modify the exciton resonance frequency in the simple model of Schmitt-Rink et al.<sup>11</sup> However, we need not do so, and the treatment in both Refs. 11 and 20 assumes a steady-state solution while the experimental conditions are more transient-like.

#### Acknowledgments

This research was supported by the Air Force Office of Scientific Research (AFSC), United States Air Force, under Contract F49620-86-C-0009, and the Office of Naval Research. The United States Government is authorized to copy and distribute reprints for governmental purposes notwithstanding any copyright notation hereon.

## References

1. R. G. Brewer and R. L. Shoemaker, Phys. Rev. Lett. 27, 631 (1971).
2. S. L. McCall and E. L. Hahn, Phys. Rev. A 2, 861 (1970).
3. R. G. Brewer, in Nonlinear Optics, ed. by P. G. Harper and B. S. Wherett (Academic, London, 1977), p. 307 ff.
4. L. Allen and J. H. Eberly, Optical Resonance and Two-level Atoms (Wiley, New York, 1975).
5. Y. R. Shen, The Principles of Nonlinear Optics (Wiley, New York, 1984).
6. A. Mysyrowicz, D. Hulin, A. Antonetti, A. Migns, W. T. Masselink and H. Morkoc, Phys. Rev. Lett. 56, 2748 (1986).
7. S. Schmitt-Rink and D. S. Chemla, Phys. Rev. Lett. 57, 2752 (1986).
8. A. Von Lenman, D. S. Chemla, J. E. Zucker and J. P. Heritage, Opt. Lett. 11, 609 (1986).
9. D. S. Chemla, D. A. B. Miller and S. Schmitt-Rink, in Optical Nonlinearities and Instabilities in Semiconductors, ed. by H. Haug (Academic, New York, 1988), p. 83 ff.
10. B. I. Greene, J. Orenstein, R. R. Millard and L. R. Williams, Chem. Phys. Lett. 139, 381 (1987); Phys. Rev. Lett. 58, 2750 (1987).
11. B. I. Green, J. F. Mueller, J. Orenstein, D. H. Rapkine, S. Schmitt-Rink and M. Thakur, Phys. Rev. Lett. 61, 325 (1988).
12. See, for example, Nonlinear Optical Properties of Organic Molecules and Crystals, ed. by D. S. Chemla and J. Zyss (Academic, New York, 1987).
13. G. M. Carter, Y. J. Chen and S. K. Tripathy, Appl. Phys. Lett. 43, 891 (1983).
14. P. Sen and P. K. Sen, IEEE J. Quantum Electron. QE23, 2033 (1987).
15. K. Singh, P. Sen and P. K. Sen, IEEE J. Quantum Electron. QE25, 67 (1989).

16. P. D. Townsend, G. L. Baker, N. E. Schlotter and S. Etemad, *Synth. Met.* 28, D633 (1989).
17. M. Thakur, Y. Shani, G. C. Chi and K. O'Brian, *Synth. Met.* 28, D595 (1989).
18. C. H. B. Cruz, J. P. Gordon, P. C. Becker, R. L. Fork and C. V. Shank, *IEEE J. Quantum Electron.* QE24, 261 (1988).
19. See D. N. Batchelder in Polydiacetylenes, ed. by D. Bloor and R. R. Chance (Martinus Nijhoff, Dordrecht, The Netherlands, 1985), p. 187 ff.
20. G. J. Blanchard, J. P. Heritage, A. C. Von Lehmen, G. L. Baker and S. Etemad, *Bull. Am. Phys. Soc.* 34, 452 (1989).
21. S. Saikan, N. Hashimoto, T. Kushida and K. Namba, *J. Chem. Phys.* 82, 5409 (1985).



# Figure captions

1. Two-level model representing the excitonic state  $|+\rangle$  and ground state  $|-\rangle$ . The virtual excitons are induced by a strong pump field with frequency  $\omega_p$ . Each transitions is mediated by a single phonon of frequency  $\omega_i$ , such that  $\omega_p + \omega_i = \omega_x$  on resonance.
2. Snapshots of the real (curve 1) and imaginary (curve 2) parts of the nonlinear susceptibility versus pump detuning at various times for  $\omega_t = \omega_x$ : (a)  $t = 25 \text{ (eV)}^{-1}$ , (b)  $t = 100 \text{ (eV)}^{-1}$ , (c)  $t = 500 \text{ (eV)}^{-1}$ , (d)  $t = 3500 \text{ (eV)}^{-1}$ .
3. Real (curve 1) and imaginary (curve 2) parts of  $\chi_t$  versus time for  $\omega_t = \omega_x$ : (a)  $\omega_p = 1.6 \text{ eV}$ ; (b)  $\omega_p = 1.72 \text{ eV}$ .
4. Snapshots of the imaginary part of the susceptibility versus probe frequency when  $\omega_p = 1.72 \text{ eV}$ : (a)  $t = 25 \text{ (eV)}^{-1}$ , (b)  $t = 500 \text{ (eV)}^{-1}$ , (c)  $t = 3500 \text{ (eV)}^{-1}$ .

Fig. 1




Fig. 2




Fig. 3





Fig. 4



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
		Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen  
Hughes Research Laboratory  
3011 Malibu Canyon Road  
Malibu, California 90265

Dr. C. B. Harris  
Department of Chemistry  
University of California  
Berkeley, California 94720

Dr. J. H. Weaver  
Department of Chemical Engineering  
and Materials Science  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. F. Kutzler  
Department of Chemistry  
Box 5055  
Tennessee Technological University  
Cookeville, Tennessee 38501

Dr. A. Reisman  
Microelectronics Center of North Carolina  
Research Triangle Park, North Carolina  
27709

Dr. D. DiLella  
Chemistry Department  
George Washington University  
Washington D.C. 20052

Dr. M. Grunze  
Laboratory for Surface Science and  
Technology  
University of Maine  
Orono, Maine 04469

Dr. R. Reeves  
Chemistry Department  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. J. Butler  
Naval Research Laboratory  
Code 6115  
Washington D.C. 20375-5000

Dr. Steven M. George  
Stanford University  
Department of Chemistry  
Stanford, CA 94305

Dr. L. Interante  
Chemistry Department  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. Mark Johnson  
Yale University  
Department of Chemistry  
New Haven, CT 06511-8118

Dr. Irvin Heard  
Chemistry and Physics Department  
Lincoln University  
Lincoln University, Pennsylvania 19352

Dr. W. Knauer  
Hughes Research Laboratory  
3011 Malibu Canyon Road  
Malibu, California 90265

Dr. K.J. Klaubunde  
Department of Chemistry  
Kansas State University  
Manhattan, Kansas 66506

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai  
Department of Chemistry  
University of California  
Berkeley, California 94720

Dr. J. Murday  
Naval Research Laboratory  
Code 6170  
Washington, D.C. 20375-5000

Dr. J. B. Hudson  
Materials Division  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. Theodore E. Madey  
Surface Chemistry Section  
Department of Commerce  
National Bureau of Standards  
Washington, D.C. 20234

Dr. J. E. Demuth  
IBM Corporation  
Thomas J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, New York 10598

Dr. M. G. Lagally  
Department of Metallurgical  
and Mining Engineering  
University of Wisconsin  
Madison, Wisconsin 53706

Dr. R. P. Van Duyne  
Chemistry Department  
Northwestern University  
Evanston, Illinois 60637

Dr. J. M. White  
Department of Chemistry  
University of Texas  
Austin, Texas 78712

Dr. D. E. Harrison  
Department of Physics  
Naval Postgraduate School  
Monterey, California 93940

Dr. R. L. Park  
Director, Center of Materials  
Research  
University of Maryland  
College Park, Maryland 20742

Dr. W. T. Peria  
Electrical Engineering Department  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson  
Department of Metallurgy and  
Materials Science  
Massachusetts Institute of Technology  
Cambridge, Massachusetts 02139

Dr. S. Sibener  
Department of Chemistry  
James Franck Institute  
5640 Ellis Avenue  
Chicago, Illinois 60637

Dr. Arnold Green  
Quantum Surface Dynamics Branch  
Code 3817  
Naval Weapons Center  
China Lake, California 93555

Dr. A. Wold  
Department of Chemistry  
Brown University  
Providence, Rhode Island 02912

Dr. S. L. Bernasek  
Department of Chemistry  
Princeton University  
Princeton, New Jersey 08544

Dr. W. Kohn  
Department of Physics  
University of California, San Diego  
La Jolla, California 92037



ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter  
Code 6170  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. Richard Colton  
Code 6170  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. Dan Pierce  
National Bureau of Standards  
Optical Physics Division  
Washington, D.C. 20234

Dr. R. Stanley Williams  
Department of Chemistry  
University of California  
Los Angeles, California 90024

Dr. R. P. Messmer  
Materials Characterization Lab.  
General Electric Company  
Schenectady, New York 12301

Dr. Robert Gomer  
Department of Chemistry  
James Franck Institute  
5640 Ellis Avenue  
Chicago, Illinois 60637

Dr. Ronald Lee  
R301  
Naval Surface Weapons Center  
White Oak  
Silver Spring, Maryland 20910

Dr. Paul Schoen  
Code 6190  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. John T. Yates  
Department of Chemistry  
University of Pittsburgh  
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene  
Code 5230  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. L. Kesmodel  
Department of Physics  
Indiana University  
Bloomington, Indiana 47403

Dr. K. C. Janda  
University of Pittsburgh  
Chemistry Building  
Pittsburgh, PA 15260

Dr. E. A. Irene  
Department of Chemistry  
University of North Carolina  
Chapel Hill, North Carolina 27514

Dr. Adam Heller  
Bell Laboratories  
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann  
Department of Chemistry  
University of Southampton  
Southampton SO9 5NH  
UNITED KINGDOM

Dr. H. Tachikawa  
Chemistry Department  
Jackson State University  
Jackson, Mississippi 39217

Dr. John W. Wilkins  
Cornell University  
Laboratory of Atomic and  
Solid State Physics  
Ithaca, New York 14853

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis  
Department of Physics  
University of California  
Irvine, California 92664

Dr. D. Ramaker  
Chemistry Department  
George Washington University  
Washington, D.C. 20052

Dr. J. C. Hemminger  
Chemistry Department  
University of California  
Irvine, California 92717

Dr. T. F. George  
Chemistry Department  
University of Rochester  
Rochester, New York 14627

Dr. G. Rubloff  
IBM  
Thomas J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, New York 10598

Dr. Horia Metiu  
Chemistry Department  
University of California  
Santa Barbara, California 93106

Dr. W. Goddard  
Department of Chemistry and Chemical  
Engineering  
California Institute of Technology  
Pasadena, California 91125

Dr. P. Hansma  
Department of Physics  
University of California  
Santa Barbara, California 93106

Dr. J. Baldeschwieler  
Department of Chemistry and  
Chemical Engineering  
California Institute of Technology  
Pasadena, California 91125

Dr. J. T. Keiser  
Department of Chemistry  
University of Richmond  
Richmond, Virginia 23173

Dr. R. W. Plummer  
Department of Physics  
University of Pennsylvania  
Philadelphia, Pennsylvania 19104

Dr. E. Yeager  
Department of Chemistry  
Case Western Reserve University  
Cleveland, Ohio 44106

Dr. N. Winograd  
Department of Chemistry  
Pennsylvania State University  
University Park, Pennsylvania 16802

Dr. Roald Hoffmann  
Department of Chemistry  
Cornell University  
Ithaca, New York 14853

Dr. A. Steckl  
Department of Electrical and  
Systems Engineering  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. G.H. Morrison  
Department of Chemistry  
Cornell University  
Ithaca, New York 14853